

Auvelity is the only NMDA receptor antagonist for MDD with oral dosing^{1,2}

Auvelity™
(dextromethorphan HBr and bupropion HCl)
extended-release tablets 45mg/105mg

One dosage strength with recommended therapeutic dose by Day 4

First 3 Days:

One tablet **once daily**

Thereafter:

One tablet **twice daily**,
taken at least 8 hours apart*

Do not take more than 2 tablets within 24 hours.

*Dosing modifications may be necessary.

Rx

Patient _____
Name _____
Address _____

AUVELITY

1 tablet PO daily x 3 days, then
increase to 1 tablet PO BID,
separated by at least 8 hours.

Dispense #60 tablets

REFILL _____ TIMES

SIGNATURE _____

DATE _____

- Should be swallowed whole and not crushed, divided, or chewed.
- Can be administered with or without food.

Before starting Auvelity:

- Assess blood pressure and monitor periodically during treatment.
- Screen for personal or family history of bipolar disorder, mania, or hypomania.
- Screen for current bupropion or dextromethorphan use in other medications.
- For patients with moderate renal impairment (eGFR of 30–59 mL/min/1.73 m²) and those who are CYP2D6 poor metabolizers: One tablet by mouth once daily in the morning.

National Drug Code (NDC): 81968-045-30

Resources are available to support you and your patients and simplify access to Auvelity. Visit [Auvelity.com/Support](https://www.auvelity.com/Support) or scan this QR code on any mobile device to learn more.



CYP2D6=cytochrome P450 2D6; eGFR=estimated glomerular filtration rate; NMDA=N-methyl-D-aspartate

INDICATION

Auvelity is indicated for the treatment of major depressive disorder (MDD) in adults.

IMPORTANT SAFETY INFORMATION

WARNING: SUICIDAL THOUGHTS AND BEHAVIORS

- Antidepressants increased the risk of suicidal thoughts and behaviors in pediatric and young adult patients in short-term studies.
- Closely monitor all antidepressant-treated patients for clinical worsening, and emergence of suicidal thoughts and behaviors.
- Auvelity is not approved for use in pediatric patients.

CONTRAINDICATIONS

Seizure: Do not use Auvelity in patients with a seizure disorder.

Current or prior diagnosis of bulimia or anorexia nervosa: A higher incidence of seizure was observed in such patients treated with bupropion.

Undergoing abrupt discontinuation of alcohol, benzodiazepines, barbiturates, and antiepileptic drugs: Due to risk of seizure.

Please see additional Important Safety Information and full [Prescribing Information](#), including **Boxed Warning** for suicidal thoughts and behaviors.

IMPORTANT SAFETY INFORMATION (CONT'D)

CONTRAINDICATIONS

Monoamine Oxidase Inhibitors (MAOIs): Do not use Auvelity concomitantly with, or within 14 days of stopping, an MAOI due to the risk of serious and possibly fatal drug interactions, including hypertensive crisis and serotonin syndrome. Conversely, at least 14 days must be allowed after stopping Auvelity before starting an MAOI antidepressant. Do not use Auvelity with reversible MAOIs such as linezolid or intravenous methylene blue.

Hypersensitivity: Do not use in patients with known hypersensitivity to dextromethorphan, bupropion, or any component of Auvelity. Anaphylactoid/anaphylactic reactions and Stevens-Johnson syndrome have been reported with bupropion. Arthralgia, myalgia, fever with rash, and other serum sickness-like symptoms suggestive of delayed hypersensitivity have also been reported with bupropion.

WARNINGS AND PRECAUTIONS

Suicidal Thoughts and Behaviors in Pediatrics and Young

Adults: Monitor all antidepressant-treated patients for any indication for clinical worsening and emergence of suicidal thoughts and behaviors, especially during the initial few months of drug therapy, and at times of dosage changes. Counsel family members or caregivers of patients to monitor for changes in behavior and to alert the healthcare provider. Consider changing the therapeutic regimen, including possibly discontinuing Auvelity, in patients whose depression is persistently worse, or who are experiencing emergent suicidal thoughts or behaviors.

Seizure: Bupropion, a component of Auvelity, can cause seizure and the risk is dose related. Because the risk of seizure with bupropion is dose-related, screen patients for use of other bupropion-containing products prior to initiating Auvelity. If concomitant use of Auvelity with other bupropion-containing products is clinically warranted, inform patients of the risk. Discontinue Auvelity and do not restart treatment if the patient experiences a seizure.

Increased Blood Pressure and Hypertension: Treatment with bupropion, a component of Auvelity, can cause elevated blood pressure and hypertension. The risk of hypertension is increased if Auvelity is used concomitantly with MAOIs or other drugs that increase dopaminergic or noradrenergic activity. Assess blood pressure before initiating treatment with Auvelity

and monitor periodically during treatment. Monitor blood pressure, particularly in patients who receive the combination of bupropion and are receiving nicotine replacement.

Activation of Mania/Hypomania: Antidepressant treatment can precipitate a manic, mixed, or hypomanic episode. The risk appears to be increased in patients with bipolar disorder or who have risk factors for bipolar disorder. Prior to initiating Auvelity, screen patients for a history of bipolar disorder and the presence of risk factors for bipolar disorder (e.g., family history of bipolar disorder, suicide, or depression). Auvelity is not approved for use in treating bipolar depression.

Psychosis and Other Neuropsychiatric Reactions: Auvelity contains bupropion and dextromethorphan. Depressed patients treated with bupropion have had a variety of neuropsychiatric signs and symptoms, including delusions, hallucinations, psychosis, concentration disturbance, paranoia, and confusion. In some cases, these symptoms abated upon dose reduction and/or withdrawal of treatment. Dextromethorphan overdose can cause toxic psychosis, stupor, coma, and hyperexcitability.

Because the risks of neuropsychiatric reactions are dose-related, screen patients for use of other bupropion- or dextromethorphan-containing products prior to initiating Auvelity. If concomitant use of Auvelity with other bupropion- or dextromethorphan-containing products is clinically warranted, monitor patients for neuropsychiatric reactions and instruct patients to contact a healthcare provider if such reactions occur.

Angle-Closure Glaucoma: The pupillary dilation that occurs following use of many antidepressants, including Auvelity, may trigger an angle closure attack in a patient with anatomically narrow angles who does not have a patent iridectomy. Avoid use of antidepressants, including Auvelity, in patients with untreated anatomically narrow angles.

Dizziness: Auvelity may cause dizziness. Precautions to reduce the risk of falls should be taken, particularly for patients with motor impairment affecting gait or a history of falls. Caution patients about operating hazardous machinery, including motor vehicles, until they are reasonably certain that Auvelity therapy does not affect them adversely.

Serotonin Syndrome: Auvelity contains dextromethorphan. Concomitant use with selective serotonin reuptake inhibitors (SSRIs) or tricyclic antidepressants increases the risk of serotonin syndrome, a potentially life-threatening condition. Prior to initiating therapy with Auvelity, screen patients for use of other dextromethorphan-containing products. If concomitant use of Auvelity with other serotonergic drugs is clinically warranted, inform patients of the increased risk for serotonin syndrome, and monitor for symptoms. Discontinue Auvelity and/or concomitant serotonergic drug(s) immediately if symptoms of serotonin syndrome occur and initiate supportive symptomatic treatment.

Embryo-fetal Toxicity: Based on animal studies, Auvelity may cause fetal harm when administered during pregnancy. Discontinue treatment in pregnant females and advise the patient about the potential risk to a fetus. Use alternative treatment for females who are planning to become pregnant.

DRUG INTERACTIONS

Strong Inhibitors of CYP2D6: Concomitant use with Auvelity increases plasma concentrations of dextromethorphan. Dosage adjustment is necessary. Monitor patients for adverse reactions potentially attributable to dextromethorphan, such as somnolence and dizziness.

Strong CYP2B6 Inducers: Concomitant use with Auvelity decreases plasma concentrations of dextromethorphan and bupropion and may decrease efficacy of Auvelity. Avoid co-administration of Auvelity.

CYP2D6 Substrates: Concomitant use with Auvelity can increase the exposures of drugs that are substrates of CYP2D6. It may be necessary to decrease the dose of CYP2D6 substrates, particularly for drugs with a narrow therapeutic index.

Digoxin: Concomitant use with Auvelity may decrease plasma digoxin levels. Monitor plasma digoxin levels in patients treated concomitantly with Auvelity.

Drugs that Lower Seizure Threshold: Concomitant use with Auvelity may increase risk of seizure. Use Auvelity with caution. Discontinue Auvelity and do not restart treatment if the patient experiences a seizure.

 **Auvelity™**
(dextromethorphan HBr and bupropion HCl)
extended-release tablets 45mg/105mg

Dopaminergic Drugs: Concomitant use with Auvelity can result in central nervous system toxicity. Use Auvelity with caution.

USE IN SPECIFIC POPULATIONS

Lactation: Because of the potential for neurotoxicity, advise patients that breast-feeding is not recommended during treatment with Auvelity and for 5 days following final dose.

Renal Impairment: Dosage adjustment is recommended in patients with moderate renal impairment (eGFR 30 to 59 mL/minute/1.73 m²). Auvelity is not recommended in patients with severe renal impairment (eGFR 15 to 29 mL/minute/1.73 m²).

Hepatic Impairment: Auvelity is not recommended in patients with severe hepatic impairment.

ADVERSE REACTIONS

Most common adverse reactions (≥5% and twice the rate of placebo): dizziness (16%), headache (8%), diarrhea (7%), somnolence (7%), dry mouth (6%), sexual dysfunction (6%), and hyperhidrosis (5%).

Please see full [Prescribing Information](#), including **Boxed Warning for suicidal thoughts and behaviors**.

AUV HCP ISI 08/2022

References: 1. Auvelity [Prescribing Information]. Axsome Therapeutics, Inc.: New York, NY. 2. FDA Depression Medicines. <https://www.fda.gov/media/132665/download>. Accessed March 21, 2022.

Please visit [Axsome.com/Disclosures](https://www.axsome.com/Disclosures) for applicable state-specific disclosures.

axsome

Auvelity, AXSOME, and its logos are trademarks or registered trademarks of Axsome Therapeutics, Inc. or its affiliates. Other trademarks are property of their respective owners.

© 2022 Axsome Therapeutics, Inc. All rights reserved.
This piece is intended for U.S. healthcare professionals.
PP-AUV-US-2200013 10/2022